anti-human Glypican - 3

4H3 Monoclonal antibody to human Glypican-3 (GPC3)

Data Sheet

Catalog # -B0038R

BACKGROUND

Glypican-3 (GPC3) is a member of the glypican family of glycosyl phosphatidylinositol-anchored cell-surface heparan sulfate proteoglycans. The 1G12 monoclonal antibody has been used to assess GPC3 expression in malignant and nonmalignant liver tissue samples and for enzyme-linked immunosorbent assay (ELISA) for detection of GPC3 in the serum. Studies have shown that GPC3 is expressed at the protein level in most HCCs, but it is undetectable in normal liver and benign hepatic lesions, including dysplastic and cirrhotic nodules. In addition, GPC3 is significantly elevated in the serum of a large proportion of patients with HCC. Based on these results, it has been proposed that GPC3 could be a useful marker to differentiate between benign and malignant liver diseases.

IMMUNOGEN

Balb/C mice immunized with the N-terminal region of the human GPC3 core protein.

SHIP CONDITIONS

Ship at ambient temperature with cold packs

STORAGE

Product should be stored at 2-8°C.

STABILITY

1 year from ship date.

REACTIVITY

Human, mouse.

REFERENCES

Capurro, M., Wanless, I. R., Sherman, M., Deboer, G., Shi, W., Miyoshi, E., and Filmus, J. Glypican-3: a novel serum and histochemical marker for hepatocellular carcinoma. Gastroenterology, 125: 89-97, 2003.

Wang X.Y., Degos F., Dubois S., Tessiore S., Allegretta M., Guttmann R.D., Jothy S., Belghiti J., Bedossa P., Paradis V. Glypican-3 expression in hepatocellular tumors: diagnostic value for preneoplastic lesions and hepatocellular carcinomas. Human Pathology. Nov; 37 (11):1435-41, 2006.

Libbrecht L., Severi T., Cassiman D., Borght S.V., Pirenne J., Nevens F., Verslype C., van Pelt J., Roskams T. Glypican-3 expression distinguishes small hepatocellular carcinomas from cirrhosis, dysplastic nodules, and focal nodular hyperplasia-like nodules. Am J Surg Pathol. Nov; 30(11):1405-11, 2006.

Di Tommaso L, Franchi G, Park YN, Fiamengo B, Destro A, Morenghi E, Montorsi M, Torzilli G, Tommasini M, Terracciano L, Tornillo L, Vecchione R and Roncalli M. Diagnostic value of HSP70, glypican 3, and glutamine synthetase in hepatocellular nodules in cirrhosis, Mar; 45 (3): 725-34, 2007.

Ordering Information

Catalog #s -B0038R Size -100 microgram Volume / vial 0.5 ml Protein G-Purified Form -Host/Clone -Mouse/4H3 Formulation - Phosphate buffered saline with 0.08% sodium azide Concentration - 0.5 milligram / ml Isotype lgG1 Application - ELISA, Western Blot, **Immunohistochemistry**